# TSE Field Trauma Triage Guidelines

## **Priority 1** (Focus: Physiology)

#### Airway/Breathing

· Actual or potential airway compromise:

Acute Hypoxia

RR < 10 or > 29 (adult)

Severe maxillofacial injuries

- Intubated/Supraglottic Airway/BVM
- Suspected inhalation injury

#### **Event/Injuries/Findings**

- Penetrating injury to head, neck, or torso
- · High voltage electrical injury
- Bilateral femur fractures
- · Complete amputation above the wrist or ankle
- Open skull fracture
- Child: Flail chest/Pelvic fracture/Pulseless extremity

#### Circulation

- · CPR by medical provider
- Hypotension:

*Adult*: SBP < 90mmHg, HR > 130

the child: Age 1-9yrs: SBP ≤ 70 + (2x age in years)

Age < 1yr: SPB ≤ 70mmHg

† Child HR: 0-12mths: > 180 or < 80bpm

≥ 12mths-5yrs: > 160 or < 60bpm

6-10yrs: > 140 or < 60bpm

≥ 11yrs: > 120 or < 60bpm

Any patient receiving blood/vasopressors

#### **Disability**

- . GCS ≤ 12 attributable to trauma
- Bilateral extremity paralysis or suspected spinal cord injury

### **Priority 2** (Focus: Anatomy)

#### Event/Injuries/Findings

- Penetrating injury proximal to elbow/knee
- Unilateral motor deficit
- · 2 or more broken extremities (any)
- Application of a tourniquet
- Open or displaced pelvic fracture
- · Open femur or humerus fracture
- Crushed or mangled extremity
- · Flail chest and/or palpable crepitus
- · Burn involvement of face, airway, hands,

feet, genitalia; OR

Adult: ≥ 20% TBSA

Child: ≥ 10% TBSA

- Pregnant(≥ 20wks) with vaginal bleeding
- Submersion with traumatic mechanism

## \*Pediatric is considered age ≤ 14yrs

## **Priority 3** (Focus: Mechanism)

#### Event/Injuries/Findings

- · Penetrating injury distal to elbow/knee
- Closed isolated femur fracture
- Loss of consciousness after injury
- GCS 13-14 after injury
- Pregnant(≥ 20wks) without vaginal bleeding
- Burn: Adult: ≤ 20% TBSA
- **†** Child: ≤ 10% TBSA Amputation of one or more digits
- Sensory deficit of an extremity

#### **Mechanisms**

Motor vehicle crash with

Death of co-occupant

Broken/bent steering wheel

Rollover

Extrication time > 20 minutes

> 12" intrusion into occupant space

Non-enclosed transport accident > 20mph

- Ejection from enclosed vehicle
- · Motor vehicle vs. pedestrian/bike
- Fall 2x patient's height
- Significant animal-related injury

#### **Consider UPGRADING one level if:**

- Pediatric: age ≤ 14yrs
- Adult: age ≥ 65yrs
- Significant co-morbidities

- Anti-coagulation other than Aspirin
- Hypothermia/Hyperthermia
- · EMS DISCRETION



Version 4, Nov 2019